# Year 9 EXPAND AND SIMPLIFY BRACKETS

2) (a) 6(m + 2t) (b) 3(3t - p)

# **Key Concepts**

#### **Expanding brackets**

Multiply the number outside the brackets with EVERY term inside the brackets

#### **Factoring expressions**

Take the highest common factor outside the bracket.

**Key Words** 

Expand

Factorise

Simplify

A hegartymaths 160, 161, 168, 189, 105, 106

	Expand and simplify where appropriate 1) $7(3 + a) = 21 + 7a$	
	2) $2(5 + a) + 3(2 + a) = 10 + 16$	2a + 6 + 3a = 5a
	3) Factorise $9x + 18 = 9(x + 2)$	
2	4) Factorise $6e^2 - 3e = 3e(2e - 1)$	
	Questions	
	<ul> <li><b>1) Expand and simplify</b></li> <li>(a) 3(2 - 7f)</li> <li>(b) 5(m - 2) + 6</li> </ul>	(c) 3(4 + t) + 2(5 + t)
	<b>2) Factorise</b> (a) 6m + 12t (b) 9t – 3p 2d	(c) 4d² –
		(c) 5q(5q – J)

(p) 2m - 4 (c) 22 + 5t

ANSWERS: 1) (a) 6 – 21f

# Year 9 **EXPANDING AND FACTORISING**

#### **Key Concepts**

**Expanding brackets** Where every term inside each bracket is multiplied by every term all other brackets.

#### **Factorising expressions**

Putting an expression back into brackets. To "factorise fully" means take out the HCF.

#### **Difference of two squares** When two brackets are

repeated with the exception of a sign change. All numbers in the original expression will be square numbers.

& hegartymaths 160-166, 168, 169,223-228

		racionse runy.
Expand and sim	plify:	1) $16at^2 + 12at = 4at(4t + 3)$
1) $4(m+5) + 3$ = $4m + 20 + 3$ = $4m + 23$ 2) $(p+2)(2p-1)$ = $p^2 + 4p - p - 2$ = $p^2 + 3p - 2$	3) $(p+3)(p-1)(p+4)$ $= (p^2+3p-p-3)(p+4)$ $= (p^2+2p-3)(p+4)$ $= 2 = p^3 + 4p^2 + 2p^2 + 8p - 3p - 12$ $= p^3 + 6p^2 + 5p - 12$	2) $x^{2} - 2x - 3 = (x - 3)(x + 1)$ 3) $6x^{2} + 13x + 5$ = $6x^{2} + 3x + 10x + 5$ = 3x(2x + 1) + 5(2x + 1) = $(3x + 5)(2x + 1)$ 4) $4x^{2} - 25$ = $(2x + 5)(2x - 5)$
<b>Key Words</b> Expand Factorise fully Bracket Difference of two squares <b>Key Words</b> A)Expand: 1) $5(m-2) + 6$ 2) $(5g-4)(2g+1)$ 3) $(y+1)(y-2)(y+3)$ B) Factorise: 1) $5b^2c - 10bc$ 2) $x^2 - 8x + 15$ 3) $3x^2 + 8x + 4$ 4) $9x^2 - 25$ $(5-x\epsilon)(5+x\epsilon)(t+(2+x)(2+x\epsilon))(\epsilon)(5-x)(2-x)(2-4))(2-4)$		

**Examples** 

Factorise fully:

# Year 9 REARRANGE AND SOLVE EQUATIONS

## **Key Concepts**

**Solving equations:** Working with inverse operations to find the value of a variable.


**Rearranging an equation:** Working with inverse operations to isolate a highlighted variable.

In solving and rearranging we **undo the operations** starting from the last one.

A hegartymaths 177-186, 280-284, 287

DELTA

Academies Trust



# Year 9 EQUATIONS IN CONTEXT

## **Key Concepts**


Algebra can be used to support us to find unknowns in a contextual problem.

We can always apply a letter to an unknown quantity, to then set up an equation.

It will often be used in area and perimeter problems and angle problems in geometry.

DELTA

Academies Trust



 $\Delta = ANNE$ ,  $L = 3 \text{ therefore the longest length is 14 cm 2) lack = 9, <math>Jane = 21$ , Sarah = 6

# Year 9 **REARRANGE AND SOLVE EQUATIONS**

## **Key Concepts**

Solving equations: Working with inverse operations to find the value of a variable.

**Rearranging an equation:** Working with inverse operations to isolate a highlighted variable.

In solving and rearranging we undo the operations starting from the last one.

& hegartymaths 177-186, 287

DELTA

Academies Trust

# Examples

Solve: **Rearrange** to make *r* the **Rearrange** to make *c* the 7p - 5 = 3p + 3subject of the formulae : subject of the formulae : -3p  $Q = \frac{2r-7}{2}$ -3p 2(3a-c) = 5c + 14p - 5 = 3expand ×3 +5 +5 6a - 2c = 5c + 1× 3 4p = 8 3Q = 2r - 7+2*c* +2*c* ÷ 2  $\div 2$ 6a = 7c + 1+7 +7 p = 2 -1 -1 3Q + 7 = 2rSolve: 6a - 1 = 7c÷ 2 ÷ 2 5(x-3) = 4(x + 2)÷ 7 ÷7  $\frac{3Q+7}{2} = r$ expand expand  $\frac{6a-1}{7} = c$ 5x - 15 = 4x + 8**-4***x* -4xx - 15 = 8+15 +15*x* = 23 Key Words 1) Solve 7(x + 2) = 5(x + 4)Solve 2) Solve 4(2 - x) = 5(x - 2)Rearrange 3) Rearrange to make m the subject 2(2p + m) = 3 - 5mTerm Inverse 4) Rearrange to make x the subject 5(x-3) = y(4-3x)Links ANSWERS: 1) x = 3 (2) x = 2 (3)  $m = \frac{7}{7}$  (4)  $x = \frac{7}{7}$  (5)  $x = \frac{7}{7}$  (7)  $x = \frac{7}{7}$ Science